



ELSEVIER



<http://www.elsevier.com/locate/jiph>

# Food animal transport: A potential source of community exposures to health hazards from industrial farming (CAFOs)

Ana M. Rule\*, Sean L. Evans, Ellen K. Silbergeld

Johns Hopkins Bloomberg School of Public Health, Department of Environmental Health Sciences, Division of Environmental Health Engineering, Baltimore, MD 21205 USA

Received 1 August 2008; accepted 13 August 2008

## KEYWORDS

Antimicrobial resistance;  
CAFO;  
Bioaerosol;  
Food animal transport;  
Air sampling;  
Surface sampling

**Summary** Use of antimicrobial feed additives in food animal production is associated with selection for drug resistance in bacterial pathogens, which can then be released into the environment through occupational exposures, high volume ventilation of animal houses, and land application of animal wastes. We tested the hypothesis that *current methods of transporting food animals from farms to slaughterhouses may result in pathogen releases and potential exposures of persons in vehicles traveling on the same road*. Air and surface samples were taken from cars driving behind poultry trucks for 17 miles. Air conditioners and fans were turned off and windows fully opened. Background and blank samples were used for quality control. Samples were analyzed for susceptible and drug-resistant strains. Results indicate an increase in the number of total aerobic bacteria including both susceptible and drug-resistant enterococci isolated from air and surface samples, and suggest that food animal transport in open crates introduces a novel route of exposure to harmful microorganisms and may disseminate these pathogens into the general environment. These findings support the need for further exposure characterization, and attention to improving methods of food animal transport, especially in highly trafficked regions of high density farming such as the Delmarva Peninsula. © 2008 King Saud Bin Abdulaziz University for Health Sciences. Published by Elsevier Ltd. All rights reserved.

**Abbreviations:** IFAP, industrial food animal production; CAFO, concentrated (or confined) animal feeding operation; ARB, antimicrobial-resistant bacteria; PTV, poultry transport vehicles; AHB, aerobic heterotrophic bacteria; PBS, phosphate buffered saline; LOD, limit of detection; cfu, colony forming units.

\* Corresponding author at: Johns Hopkins Bloomberg School of Public Health, Department of Environmental Health Sciences, Division of Environmental Health Engineering, 615 N Wolfe Street, Room E6618, Baltimore, MD 21205, USA. Tel.: +1 410 502 5952; fax: +1 410 955 9334.

E-mail address: [arule@jhsph.edu](mailto:arule@jhsph.edu) (A.M. Rule).

## Introduction

The industrial food animal production (IFAP) environment, of which some operations (depending upon size) are also known as concentrated animal feeding operations (CAFOs), is a well characterized source of antibiotic-resistant pathogenic bacteria, with documented exposures to human populations via multiple pathways [1–4]. Our research in this area has characterized both food and environmental pathways of exposure to antimicrobial-resistant bacteria (ARB) in the context of broiler poultry production on the Delmarva Peninsula, one of the major production areas in the US [5–7].

The purpose of this project was to investigate a hitherto unstudied route of potential community exposures to ARBs from poultry production: the transport of live flocks from poultry houses to processing plants. Food animals are customarily transported from CAFOs to slaughterhouses in pens, open cages, or crates stacked on flat bed trailers or in a trailer with little or no containment. These methods provide no barrier to pathogens into the environment. Transport cages are known to become highly contaminated with feces and bacteria during the process of transport [8–13]. Ramabu et al. [14], in a study of *Campylobacter* contamination within the broiler poultry production environment, found that after transport, poultry crates were contaminated at a rate of 75%, while 47% of truck beds were also contaminated. This phenomenon has been associated with the contamination of flocks as they enter the processing plant [15–17].

This potential route of environmental release is of public health importance to both local and regional communities. In the Delmarva Peninsula, the opportunities for broader population exposure are considerable, since during the summer months hundreds of thousands of vehicles cross through the same roads trafficked by poultry transport vehicles (PTVs) in order to reach major recreation areas in all three states. *The specific aim of this project was to test the hypothesis that persons riding in passenger cars behind live broiler PTVs are at risk of exposure to antibiotic-resistant bacteria originating from uncontrolled releases of feces and other materials from the open crates of PTV.*

## Methods

### Feasibility study

Prior to undertaking this study, we first validated methods of sample collection in cars through trial runs in Baltimore City and in the Delaware, Mary-

land and Virginia (Delmarva) peninsula. We ran several experimental runs in the same or nearby locations as the routes utilized by PTVs. After confirming methods for air and surface sampling, we conducted three environmental sampling trips, from June through October 2007.

### Sampling site and collection

This study was conducted on the Delmarva Peninsula, a region with the highest density of broiler chickens per farm acre in the United States, in which several major poultry processing plants are also located [18] (see Fig. 1; area of sampling is indicated by the dashed oval). The route of sampling (route 13) was selected because it is the only access road from farms in Maryland to processing plants in Virginia. We chose PTVs traveling to a processing plant in Accomack, VA, since this facility was located approximately 17 miles (~20 min) south from the start point for this study, which



Figure 1 Delmarva peninsula, with the sampling area indicated by the dashed oval. Map of the Delmarva Peninsula provided by: [www.worldatlas.com](http://www.worldatlas.com).

maximized the time for following each PTV. Each sampling run was carried out by a four door passenger vehicle traveling from the starting point to the plant. Briefly, each vehicle followed a truck (2–3 car lengths distance behind) for 17 miles until reaching the processing plant. Air conditioners and fans were turned off and all four windows were fully opened during each sampling run. Three separate air sampling events were conducted on June 20 2007, August 14 2007, and October 18 2007, consisting of 1 run, 6 runs, and 3 runs respectively, for a total of 10 runs. Surface sampling was conducted only on the August and October runs, for a total of 5 runs. Start and end times, temperature and relative humidity were recorded for each run.

Air samples were collected using two methods: a sterile all-glass sampler (BioSampler®, SKC Inc) with 20 ml sterile 1× PBS as collection media and 25 mm gelatin filters (Sartorius, Germany) on sterile open-faced cassettes (Derlin 1107, Pall Life Sciences, Ann Arbor, MI). The BioSampler® collected air samples at a calibrated flow rate of 12.5 L/min, while the open-faced filter had a calibrated flow rate of 4 L/min. The BioSampler was placed in the center of the back seat, with the air intake facing directly to the front. Air was drawn with a vacuum pump (VP0435A, MEDO USA) connected through a power inverter (Pwrinv 400 W, AIMS Corp., Taiwan, ROC) to the vehicle's power. The filter holder was clipped to the side of the driver's head rest at a slightly downward angle (to simulate normal breathing and avoid direct impaction). Both samplers are designed to collect bioaerosols larger than 0.3 μm in diameter. At the termination of each run (within 10 min) the filters were aseptically placed in tubes containing sterile PBS, and the remaining BioSampler® volumes (the air flow caused some evaporation) were recorded and pipetted into sterile tubes.

Surface samples were collected using sterile swabs, wetted with sterile PBS, from two designated surfaces: external, from the outside door handle on the driver's side, and internal: from the top surface of an unopened soda can placed in the cup holder between the driver and front passenger seats. Wipes were collected by wiping for approximately 20 s, and then stirred into sterile PBS in a tube for another 20–30 s.

Blank and background control samples were also collected for both air and surface sampling. Blank air control samples consisted of collection liquid and filters that were inside the car but not exposed to the air. Surface blanks were collected immediately after swiping the target surface with an alcohol swab. Background control air samples were collected during a drive in the same road, in the

absence of a PTV, using the same sampling protocol as described above; surface background controls were collected after this control drive. All samples were stored on ice, and processed immediately upon returning to the laboratory (within 8 h).

### Bacterial isolation and enumeration

Total aerobic heterotrophic bacteria (AHB) and *Enterococcus* spp. samples were isolated and enumerated using respective agar media, all obtained from Becton Dickinson (Sparks, MD). Samples were analyzed for culturability using standard dilution and spread-plating methods, as well as membrane filtration technique [19]. Briefly, a 1 mL aliquot of each sample was passed through a 47 mm diameter, 0.45 μm pore size cellulose membrane filter (GE Water & Processing Technologies) using a filter funnel and vacuum system. Microorganisms present in the sample are retained on the surface of the filter which is then placed on a nutrient agar plate. In addition, 100 μL aliquots of each sample are plated on nutrient agar. AHB spp. were isolated on tryptic Soy agar, while *Enterococcus* spp. were isolated on enterococcosel agar. All samples were plated in triplicate. Tryptic soy agar plates were incubated at 37 °C for 24 h, while enterococcosel agar plates were incubated at 37 °C for 36 h. Blank and background controls were included in all analyses.

Resulting colonies were counted, and used to determine the concentration of isolated bacteria per cubic meter of air or per square centimeter of surface area as per Eqs. (1) and (2). Enterococcal colonies were indicated by the formation of a black precipitate on enterococcosel agar. Selected presumptive enterococci isolates (Note: previous research on air samples [20] showed that not all isolates growing on this media were *Enterococcus* spp., and therefore we will use the term "presumptive") were re-plated on enterococcosel agar and archived in 10% glycerol–tryptic soy broth at –80 °C for subsequent antimicrobial susceptibility testing.

$$\frac{\text{cfu}}{\text{m}^3_{\text{air}}} = \frac{\text{cfu}}{V_p (\text{mL}_{\text{liq}})} \times \frac{V_f (\text{mL}_{\text{liq}})}{Q_s (\text{L}_{\text{air}}/\text{min}) \times t_s (\text{min})} \times \frac{1000 (\text{L}_{\text{air}})}{\text{m}^3_{\text{air}}} \quad (1)$$

$$\frac{\text{cfu}}{\text{cm}^2} = \frac{\text{cfu}}{V_p (\text{mL}_{\text{liq}})} \times \frac{V_f (\text{mL}_{\text{liq}})}{A_s (\text{cm}^2)} \quad (2)$$

where cfu is the colony forming units per plate,  $V_p$  the volume of liquid plated ( $\text{mL}_{\text{liq}}$ ),  $V_f$  the final volume of liquid from where samples were plated

(mL<sub>liq</sub>),  $Q_s$  the air flow rate (L<sub>air</sub>/min),  $T_s$  the sampling time (min), and  $A_s$  is the area swiped (cm<sup>2</sup>).

Due to poor recovery of resistant isolates during enterococcal isolation of samples collected from runs 1 and 2 using the protocol mentioned above, we decided to increase the sensitivity of the assay for detection of resistant enterococcal isolates in the samples from run 3 by enriching samples in nutrient broth supplemented with antibiotics at breakpoint concentrations [5]. This strategy optimizes the likelihood of recovering resistant isolates if these are present. Isolates were enriched in enterococcosel broth with the following antibiotics (concentrations): tetracycline (8 µg/ml), vancomycin (16 µg/ml), erythromycin (4 µg/ml), and quinupristin–dalbopristin (2 µg/ml).

Since most of the background samples were below the limit of detection (<LOD), a method LOD was calculated for comparison purposes. This is determined by the minimum number of colony forming units (cfu) per plate, following EPA SW 846 [21]. Eqs. (3) and (4) show the LOD for each method, which was used for graphically representing our data, and where  $V_s$  is the volume of total sample,  $V_p$  is the volume of sample plated,  $A_s$  is the total surface area swiped, and  $V_a$  is the volume of air sampled.

$$\text{LOD surface} \left( \frac{\text{cfu}}{\text{cm}^2} \right) = \frac{5 \text{ cfu}}{V_p (\text{mL}_{\text{liq}})} \times \frac{V_s (\text{mL}_{\text{liq}})}{\text{swab}} \times \frac{1 \text{ swab}}{A_s (\text{cm}^2)} \quad (3)$$

$$\text{LOD air} \left( \frac{\text{cfu}}{\text{m}^3_{\text{air}}} \right) = \frac{5 \text{ cfu}}{V_p (\text{mL}_{\text{liq}})} \times \frac{V_s (\text{mL}_{\text{liq}})}{\text{sample}} \times \frac{\text{sample}}{V_a (\text{m}^3_{\text{air}})} \quad (4)$$

### Antibiotic susceptibility testing

The disk diffusion method [22] was used to assess the susceptibility of presumptive enterococci isolates ( $n=104$ ) to eight different antimicrobial drugs. The antimicrobial drugs (including concentrations) used to test for susceptibility were selected based on our previous studies in the broiler poultry environment: penicillin (10 IU), erythromycin (15 µg), virginiamycin [streptogramin A and B combination] (15 µg), vancomycin (30 µg), streptomycin (300 µg), gentamicin (120 µg), ciprofloxacin (5 µg), and tetracycline (30 µg). All drug-impregnated disks were obtained from Becton Dickinson (Sparks, MD).

To perform the disk diffusion test, archived enterococcal isolates were grown on Tryptic soy agar at 37 °C under aerobic conditions. After 24 h, each bacterial sample was suspended in 4 mL Mueller–Hinton broth with a sterile inoculating loop and adjusted to a 0.5 McFarland standard using a Vitek colorimeter (Hach, Loveland, CO). Using a sterile cotton swab, a standard lawn of each sample was created on Mueller–Hinton agar plates. This was followed by dispensing the antimicrobial disks on each plate providing 33 mm distance between disk centers using a disk dispenser obtained from Becton Dickinson (Sparks, MD). Plates were incubated at 37 °C for 24 h.

The zones of inhibition created by exposure to each antimicrobial drug were measured using a millimeter ruler. The following zone diameters (in mm) correspond to resistance, intermediacy, and susceptibility to respective antibiotics: penicillin [ $\leq 14$ ,  $-$ ,  $\geq 15$ ]; erythromycin [ $\leq 13$ , 14–22,  $\geq 23$ ]; virginiamycin [ $\leq 15$ , 16–18,  $\geq 19$ ]; vancomycin [ $\leq 14$ , 15–16,  $\geq 17$ ]; streptomycin [ $\leq 6$ , 7–9,  $\geq 10$ ]; gentamicin [ $\leq 6$ , 7–9,  $\geq 10$ ]; ciprofloxacin [ $\leq 15$ , 16–20,  $\geq 21$ ]; and tetracycline [ $\leq 14$ , 15–18,  $\geq 19$ ] [22].

### Statistical analysis


Statistical significance was considered at  $\alpha=0.05$  level, and tested using STATA® (College Station, TX). Differences between the air samples ( $N=9$ ) and their respective backgrounds ( $N=4$ ) were tested using two-sample  $t$ -tests with equal variances. Differences between the two air samplers were tested with paired  $t$ -tests. Differences between the swab samples ( $N=5$  for internal and 3 for external) and their respective controls ( $N=4$  for internal and 3 for external) were determined with Wilcoxon rank-sum non-parametric tests, due to small sample size.

## Results

### Enumeration of AHB and enterococci from air and surface samples

Results are presented as concentrations in comparison to background. Background for all methods was <5 cfu/plate. This count was used to establish the LOD for each method.

Based on Eqs. (3) and (4), LOD for the interior wipe = 1.25 cfu/cm<sup>2</sup>; LOD for the exterior wipe = 0.25 cfu/cm<sup>2</sup>; LOD for the BioSampler samples = 400 cfu/m<sup>3</sup>, and LOD for the filter samples = 500 cfu/m<sup>3</sup>.




**Figure 2** Increased levels of airborne AHB (aerobic heterotrophic bacteria) and presumptive enterococci (ENT) as compared to background levels from both BioSampler (BS) and gelatin filter (Fltr). Data for all 10 runs, June 20, August 14, October 18, 2007. ABkg = Air Background Control. The box extends from the 25th to the 75th percentile. The line across the box represents the median value. The whiskers extend to the most extreme observations that are within 1.5 times the percentiles. Points outside this range represent outliers. ✱: statistically significant from background.

### Air samples

Data for all 10 runs are presented in Fig. 2. Mean concentrations of airborne AHB collected while driving behind PTVs were  $2.8 \times 10^3$  cfu/m<sup>3</sup> for BioSamplers and  $3.6 \times 10^3$  cfu/m<sup>3</sup> for the filters, both statistically higher than background controls (all background controls were <LOD) ( $p < 0.05$ ). Average concentrations of presumptive airborne enterococci for the control runs were <LOD, with concentrations of  $1.3 \times 10^3$  cfu/m<sup>3</sup> from the BioSampler ( $p > 0.05$ ) and  $2.3 \times 10^3$  from the filter sampler ( $p < 0.01$ ).

### Surface samples

Interior and exterior wipe samples were positive after driving behind PTVs. Background levels for AHB and presumptive enterococci were below the limit of detection (<1.25 and <0.25 cfu/cm<sup>2</sup> for both interior and exterior surfaces respectively) on both dates. Average levels on the interior surface after driving behind PTV were 18 cfu/cm<sup>2</sup> for AHB ( $p < 0.05$ ) and 9 cfu/cm<sup>2</sup> for presumptive enterococci ( $p > 0.05$ ). Average external AHB and presumptive enterococci levels were 2.8 and 2.3 cfu/cm<sup>2</sup>, respectively, both higher than background ( $p > 0.05$ ). All samples were above the limit of detection. Data are presented in Fig. 3.




**Figure 3** Increased levels of AHB (aerobic heterotrophic bacteria) and presumptive enterococci (ENT) from surface samples as compared to background. External control (Ext ctrl) and Internal control (Int ctrl) are background controls (not following poultry transport vehicles (PTV)). Internal/External sample represent surface samples (following PTV) in cfu/cm<sup>2</sup>.  $N = 5$  for the internal samples,  $N = 4$  for internal controls, and  $N = 3$  for external controls and samples. The box extends from the 25th to the 75th percentile. The line across the box represents the median value. The whiskers extend to the most extreme observations that are within 1.5 times the percentiles. Points outside this range represent outliers. ✱: statistically significant from background; ●: data outlier.

### Antibiotic-resistant enterococci

Tetracycline, erythromycin, quinupristin/dalfopristin, and high level streptomycin resistance phenotypes were detected among isolated presumptive *Enterococcus* spp. ( $n = 24$  out of 96, 25%) recovered from airborne and surface samples after traveling behind poultry transport vehicles. No resistant isolates were detected in background samples. Of the 24 resistant isolates, 62.5% were resistant to tetracycline (Tet<sup>r</sup>), 41.7% were resistant to erythromycin (Ery<sup>r</sup>), one isolate was resistant to quinupristin/dalfopristin and one isolate was resistant to high level concentrations of streptomycin. Three of the 24 isolates were resistant to more than one antibiotic. There was no resistance to vancomycin detected within any of the isolates (Fig. 4).

### Discussion

Increased concentrations of AHB and presumptive enterococci were isolated from the air and internal/external surfaces of cars traveling behind PTVs. This confirms the possibility of major pathogen dispersal via this route, and highlights the



**Figure 4** Phenotypic profile for resistant isolates. Percentage of resistant and total isolates resistant to individual antibiotics and multiple antibiotics. Tet: tetracycline, Ery: erythromycin, Q/D: quinupristin/dalfopristin, HL Strep: high level streptomycin, Pen: penicillin, HL Gent: high level gentamicin, Cip: ciprofloxacin, Van: vancomycin, MD: multi-drug.

importance of better containment of animals and waste. Surface samples indicate that airborne bacteria, including ARB strains, can deposit on surfaces and remain viable for up to 20 min. Twelve presumptive enterococci isolates were found to be resistant to three antimicrobial drugs of human importance: tetracycline, erythromycin, and quinupristin/dalfopristin. These three drugs are all approved for use in broiler poultry [23]. These data are consistent with previous research by us and others that has demonstrated associations between antibiotic use in food animal production and the presence of microbial antibiotic resistance [3,5]. Also of importance, we detected no evidence for vancomycin resistance, consistent with other studies in the food animal environment conducted by us and by Chapin et al. [20]. Since vancomycin analogs have never been used in US food animal production [24], this finding supports the inference that the source of ARB in our samples was not likely to be from humans.

Previous reports have revealed heightened risks of ARB exposure to the public and occupational workers through consumption and handling of raw, improperly cooked or live poultry [6]. Water and air sources near and within animal facilities have also been implicated as sources of exposure to drug-resistant bacteria [20,25,26]. Currently, there are no data on air releases from poultry houses. We and others have also reported that workers handling live poultry are at increased risk of exposure to ARB [6,7]. The results from this study indicate that the current practice of transporting

poultry (and very likely other food animals) may serve as another route of exposure that requires attention. Populations at increased risk to this exposure pathway include motorists and others traveling behind PTV, as well as neighborhood populations along the PTV path, which likely include susceptible populations such as children and elderly people.

This study has also served to validate the use of gelatin filters for sampling biological aerosols inside a moving vehicle. Gelatin filter data were not statistically different ( $p > 0.1$ ) from the widely used and validated BioSampler [27]. This is a significant finding, since collection of bioaerosols onto filters presents advantages over liquid samplers because of their portability and ease of use. The use of membrane filters has traditionally been problematic due to low survival efficiency caused by cell desiccation and poor microbial recovery from the filters [28]. Gelatin filters offer a moist surface conducive to cell viability and the ability to dissolve completely within the diluent, which precludes the need for cell extraction from the filter.

This first study is limited in size and scope. Moreover, during the conduct of the runs we were aware of several factors that may contribute to variability in results. Weather, including temperature, humidity, and wind direction/velocity, may affect levels of bacteria in air and impact on cars [29]. The speed and distance between PTVs and receiver vehicles, which could not be controlled, also varied among vehicles and runs. Given the empirical nature of this sampling, it is not possible to control all these variables and thus substantially more sampling will be required to more precisely estimate risks of exposure, and to support generalization to other poultry producing areas.

Despite the abovementioned limitations, all but one of the air samples and inside surface swipes yielded statistically elevated bacteria from background, and ARB strains were isolated from both air and surface samples. These data are a strong indication that PTVs are a likely route of exposure to antibiotic-resistant bacteria. These findings support the need for further exposure characterization, and attention to improving methods of food animal transport, especially in highly trafficked regions of high density farming such as the Delmarva Peninsula.

## Acknowledgements

Support for this research was received from an Innovation grant from the Johns Hopkins Center for a Livable Future. The authors would like to thank Dr.

Jay Graham and Carol Resnick for their invaluable help.

## References

- [1] Silbergeld EK, Graham J, Price LB. Industrial food animal production, antimicrobial resistance, and human health. *Ann Rev Pub Health* 2008;29:151–69.
- [2] Singer RS, Finch R, Wegener HC, Bywater R, Walters J, Lipsitch M. Antibiotic resistance—the interplay between antibiotic use in animals and human beings. *Lancet Infect Dis* 2003;3(1):47–51.
- [3] Wegener HC. Antibiotics in animal feed and their role in resistance development. *Curr Opin Microbiol* 2003;6(5):439–45.
- [4] Hershberger E, Oprea SF, Donabedian SM, Perri M, Bozigar P, Bartlett P, et al. Epidemiology of antimicrobial resistance in enterococci of animal origin. *J Antimicrob Chemother* 2005;55(1):127–30.
- [5] Price LB, Johnson TE, Vailes R, Silbergeld E. Fluoroquinolone-resistant campylobacter isolates from conventional and antibiotic-free chicken products. *Environ Health Perspect* 2005;113(5):557–60.
- [6] Price LB, Roess A, Graham JP, Baqar S, Vailes R, Sheikh KA, et al. Neurologic symptoms and neurologic antibodies in poultry workers exposed to *Campylobacter jejuni*. *J Occup Environ Med* 2007;49(7):748–55.
- [7] Price LB, Graham JP, Lackey LG, Roess A, Vailes R, Silbergeld EK. Elevated risk of carrying gentamicin-resistant *Escherichia coli* among US poultry workers. *Environ Health Perspect* 2007;115(12):1738–42.
- [8] Berrang ME, Northcutt JK, Cason JA. Recovery of *Campylobacter* from broiler feces during extended storage of transport cages. *Poult Sci* 2004;83(7):1213–7.
- [9] Hayes JR, Wagner DD, English LL, Carr LE, Joseph SW. Distribution of streptogramin resistance determinants among *Enterococcus faecium* from a poultry production environment of the USA. *J Antimicrob Chemother* 2005;55(1):123–6.
- [10] Ramesh NS, Joseph W, Carr LE, Douglass LW, Wheaton FW. Serial disinfection with heat and chlorine to reduce microorganism populations on poultry transport containers. *J Food Protect* 2003;66(5):793–7.
- [11] Corry JE, Allen VM, Hudson WR, Breslin MF, Davies RH. Sources of *Salmonella* on broiler carcasses during transportation and processing: modes of contamination and methods of control. *J Appl Microbiol* 2002;92(3):424–32.
- [12] Slader JG, Domingue F, Jorgensen K, McAlpine RJ, Bolton Owen FJ, Humphrey TJ. Impact of transport crate reuse and of catching and processing on *Campylobacter* and *Salmonella* contamination of broiler chickens. *Appl Environ Microbiol* 2002;68(2):713–9.
- [13] Rigby CE, Pettit JR, Bentley AH, Spencer JL, Salomons MO, Lior H. The relationships of salmonellae from infected broiler flocks, transport crates or processing plants to contamination of eviscerated carcasses. *Can J Comp Med* 1982;46(3):272–8.
- [14] Ramabu SS, Boxall NS, Madie P, Fenwick SG. Some potential sources for transmission of *Campylobacter jejuni* to broiler chickens. *Lett Appl Microbiol* 2004;39(3):252–6.
- [15] Herman LM, Heyndrickx K, Grijspeerdt D, Vandekerchove I, Rollier I, De Zutter L. Routes for *Campylobacter* contamination of poultry meat: epidemiological study from hatchery to slaughterhouse. *Epidemiol Infect* 2003;131(3):1169–80.
- [16] Heyndrickx M, Vandekerchove D, Herman L, Rollier I, Grijspeerdt K, De Zutter L. Routes for *Salmonella* contamination of poultry meat: epidemiological study from hatchery to slaughterhouse. *Epidemiol Infect* 2002;129(2):253–65.
- [17] Joseph SW, Hayes JR, English LL, Carr LE, Wagner DD. Implications of multiple antimicrobial-resistant enterococci associated with the poultry environment. *Food Addit Contam* 2001;18(12):1118–23.
- [18] Ringbauer Jr JA, James JB, Genthner FJ. Effects of large-scale poultry farms on aquatic microbial communities: a molecular investigation. *J Water Health* 2006;4(1):77–86.
- [19] Eaton A. Standard methods for the examination of water and waste water. Washington, DC: APHA/AWWA/WEF; 1995.
- [20] Chapin A, Rule A, Gibson K, Buckley T, Schwab K. Airborne multidrug-resistant bacteria isolated from a concentrated swine feeding operation. *Environ Health Perspect* 2005;113(2):137–42.
- [21] US EPA. Test methods for evaluating solid waste. Laboratory manual physical/chemical methods, vol. 1A, 3rd ed. Washington, DC: US Environmental Protection Agency; 1986 [SW-846].
- [22] National Committee for Clinical Laboratory Standards. Performance standards for antimicrobial susceptibility testing. Wayne, PA: National Committee for Clinical Laboratory Standards; 2002.
- [23] Florini KR, Denison T, Stiffler T, Fitzgerald T, Goldberg R. Resistant bugs and antibiotic drugs: state and county estimates of antibiotics in agricultural feed and animal waste. USA: Environmental Defense; 2005.
- [24] Sapkota AR, Lefferts LY, McKenzie S, Walker P. What do we feed to food production animals? A review of animal feed ingredients and their potential impacts on human health. *Environ Health Perspect* 2007;115:663–70.
- [25] Gibbs SG, Green CF, Tarwater PM, Scarpino PV. Resistant and nonresistant bacteria recovered from two swine herd confined animal feeding operations. *J Occup Environ Hyg* 2004;1:699–706.
- [26] Sapkota AR, Curriero FC, Gibson KE, Schwab KJ. Antibiotic-resistant enterococci and fecal indicators in surface waters and groundwater impacted by a concentrated swine feeding operation. *Environ Health Perspect* 2007;115:1040–5.
- [27] Lin XJ, Reponen T, Willeke K, Wang Z, Grinshpun SA, Trunov M. Survival of airborne microorganisms during swirling aerosol collection. *Aerosol Sci Technol* 2000;32(3):184–96.
- [28] Rule AM, Kesavan J, Schwab KJ, Buckley TJ. Application of flow cytometry for the assessment of preservation and recovery efficiency of bioaerosol samplers spiked with pan-toea agglomerans. *Environ Sci Technol* 2007;41:2467–72.
- [29] Muilenberg ML, Skellenger WS, Burge HA, Solomon WR. Particle penetration into the automotive interior. I: Influence of vehicle speed and ventilatory mode. *J Allergy Clin Immunol* 1991;87(2):581–5.